

Simplicity of $QC(\mathbb{S}^n)$ and $LIP(\mathbb{S}^n)$

James Giblin

February 1, 2008

ABSTRACT

We will prove that the groups $QC(\mathbb{S}^n)$ and $LIP(\mathbb{S}^n)$ ($n \geq 2$) are simple.

1 Introduction

A group G is simple if its only normal subgroups are itself and the trivial group. Let $QC(\mathbb{S}^n)$ and $LIP(\mathbb{S}^n)$ denote the orientation preserving quasiconformal and bilipschitz homeomorphisms of $\mathbb{S}^n$ respectively (We will implicitly assume throughout that the dimension n is greater than 1). In this paper we prove the following statement,

Theorem 1.1 *The groups $QC(\mathbb{S}^n)$ and $LIP(\mathbb{S}^n)$ are simple.*

Our proof falls into two stages. First, we show that the subgroups of $QC(\mathbb{S}^n)$ and $LIP(\mathbb{S}^n)$ which are generated by the elements supported on open balls are simple. In the second stage, we show that *every* element of $QC(\mathbb{S}^n)$ or $LIP(\mathbb{S}^n)$ can be written as a composition of elements which are all supported on open balls. The main part of the argument is Lemma 3.6 which proves that there are neighbourhoods of the identity in $QC(\mathbb{S}^n)$ and $LIP(\mathbb{S}^n)$ which satisfy this condition.

Anderson [1] proved that if G is a group of homeomorphisms of a Hausdorff space X and K is a basis for X satisfying certain conditions, then the group G_0 generated by the homeomorphisms which are supported on elements of K is simple. In fact, he showed that if h is any non trivial element of G , then every element of G_0 can be written as a product of four conjugates of h and h^{-1} . We demonstrate in detail that the group of quasiconformal homeomorphisms of $\mathbb{S}^n$ or the group of Lipschitz homeomorphisms of $\mathbb{S}^n$ and the basis consisting of open balls satisfy the necessary conditions to apply his argument and give the proof that these groups will therefore be simple.

In his paper, Anderson also gave some conditions that the whole group G will be generated by G_0 . Namely that X can be expressed as the union of two

basis elements, and that G has the "Annulus property". Here, G satisfies the annulus property if for every $g \in G$ and $k \in K$ such that there exists $k' \in K$ with $k \cup g(k) \subset k'$, there exists $g_0 \in G_0$ such that g and g_0 agree on k . Now, $\mathbb{S}^n$ can obviously be written as the union of two open balls and in the process of proving Lemma 3.6 we show that there are neighbourhoods of id in the groups $QC(\mathbb{S}^n)$ and $LIP(\mathbb{S}^n)$ which satisfy the annulus property. In these cases, this turns out to be enough.

A homeomorphism $h : X \rightarrow X$ is called *stable* if it can be written as a finite composition of homeomorphisms, each of which is the identity on some open set. We say a group of homeomorphisms G is *stable* if every element of G is (See [3] and [4] for more information on stable homeomorphisms). Kirby [7] showed that the group of homeomorphisms of the torus $\mathbb{T}^n$ is stable. He used this, along with an immersion of $\mathbb{T}^n$ minus a point into $\mathbb{R}^n$ to show that the group of homeomorphisms of $\mathbb{R}^n$ is locally contractible. Edwards and Kirby [5] then went on to extend these results to arbitrary manifolds, in particular, showing that the homeomorphism group of a compact manifold M is locally contractible.

The stable homeomorphism conjecture says that the orientation preserving homeomorphisms of $\mathbb{R}^n$ are stable. This is known to be true for $n \neq 4$ [7] whereas for $n = 4$ this question is unsolved. A simple application of Anderson's results concludes from the stable homeomorphism conjecture that the group of orientation preserving homeomorphisms of $\mathbb{S}^n$ is simple for $n \neq 4$.

In general, if M is a non compact manifold then its group of homeomorphisms will not be simple, as the subgroup consisting of elements with compact support forms a non-trivial normal subgroup. Likewise for manifolds with boundary, the subgroup consisting of elements which act as the identity in some neighbourhood of the boundary will also be a non-trivial normal subgroup. In the case of an arbitrary closed manifold the homeomorphism group has generally got more than one path component, with the component of the identity (which consists of all the stable homeomorphisms) forming a non-trivial normal subgroup. It is still an interesting question to ask whether these subgroups are themselves simple.

For the case of diffeomorphisms, let M be an n -dimensional smooth manifold. Define $\text{Diff}(M, r)$ to be the group of C^r diffeomorphisms of M which are isotopic to the identity through compactly supported C^r isotopies (An isotopy has compact support if there is compact set outside of which the isotopy acts trivially). In [9] and [10] Mather proved that if $\infty \geq r \neq n + 1$ then $\text{Diff}(M, r)$ is perfect (equal to its own commutator subgroup) with the case $r = \infty$ being proved by Thurston (See [14]). Now, in [6], Epstein showed that for certain groups of homeomorphisms, the commutator subgroup is simple. In particular, $\text{Diff}(M, r)$ is simple when $\infty \geq r \neq n + 1$.

Simplicity questions have also been addressed in the symplectic and volume preserving categories. Perhaps most notably Banyaga [2] proved that for a closed symplectic manifold M the group of Hamiltonian symplectomorphisms $\text{Ham}(M, \omega)$ is perfect and then used Epstein's argument, which also applies in the symplectic case, to deduce that it is simple. On a non compact manifold, if one considers the component of the identity in the group of compactly supported symplectomorphisms $\text{Symp}_0^c(M, \omega)$, then there exists a surjective homomorphism from $\text{Symp}_0^c(M, \omega)$ to $\mathbb{R}$. Banyaga proved that the kernel of this homomorphism is simple and also showed an analogous result in the volume preserving case. In general however, little is known about the normal subgroups of $\text{Symp}(M, \omega)$. For example we currently do not know whether the group of symplectomorphisms of $\mathbb{R}^{2n}$ with its standard symplectic form is perfect. A stark contrast to the volume preserving case, where McDuff [11] has shown that $\text{Diff}_{\text{vol}}(\mathbb{R}^n)$ is perfect for every dimension $n \geq 3$.

In [12] Sullivan proved the stable homeomorphism conjecture and the annulus conjecture in the locally quasiconformal and Lipschitz categories in all dimensions. We go through his proof of the stable homeomorphism conjecture in detail, and use the result to show simplicity of $\text{QC}(\mathbb{S}^n)$ and $\text{LIP}(\mathbb{S}^n)$.

I would like to thank Vladimir Markovic for introducing me to this problem and for many helpful discussions regarding this work.

2 QC and LIP homeomorphisms supported on open balls generate a simple group.

In this section we start by introducing some notation and terminology and go on to prove that every element of $\text{QC}(\mathbb{S}^n)$ or $\text{LIP}(\mathbb{S}^n)$ which is supported on an open ball can be written as a composition of four conjugates of an arbitrary non-trivial element and its inverse. More importantly for us, this gives us the easy corollary that the subgroups of $\text{QC}(\mathbb{S}^n)$ and $\text{LIP}(\mathbb{S}^n)$ generated by those elements supported on open balls are simple.

Let $d_{\mathbb{R}^n}$ denote the usual Euclidean distance on $\mathbb{R}^n$, we set

$$\mathbb{S}^{n-1} = \{x \in \mathbb{R}^n : d_{\mathbb{R}^n}(x, 0) = 1\}$$

to be the unit sphere in $\mathbb{R}^n$ and denote the standard spherical metric on $\mathbb{S}^n$ by $d_{\mathbb{S}^n}$. Let

$$\mathbb{B}^n = \{x \in \mathbb{R}^n : d_{\mathbb{R}^n}(x, 0) < 1\}$$

denote the unit ball in $\mathbb{R}^n$, and let $d_{\mathbb{B}^n}$ denote its hyperbolic metric. We now define,

$$B_X(c, \epsilon) = \{x \in X : d_X(x, c) < \epsilon\}$$

to be the open ball of radius $\epsilon > 0$ around $c \in X$, where X is one of $\mathbb{R}^n$, $\mathbb{S}^n$ or $\mathbb{B}^n$ and d_X is the appropriate metric.

We also define a metric on the space of maps from $\mathbb{S}^n$ to itself by,

$$\tilde{d}(f, g) = \sup\{d(f(x), g(x)) : x \in \mathbb{S}^n\}.$$

The group of Möbius transformations $\text{Möb}(\mathbb{S}^n)$ is the transformation group of $\mathbb{S}^n$ generated by inversions in $(n - 1)$ -spheres and $\text{Möb}^+(\mathbb{S}^n)$ is the index two subgroup consisting of those which preserve orientation. By identifying $\mathbb{S}^n$ with $\mathbb{R}^n \cup \{\infty\}$ we can think of $\mathbb{B}^n$ as a subset of $\mathbb{S}^n$. If we define $\text{Möb}(\mathbb{B}^n) = \{f \in \text{Möb}(\mathbb{S}^n) : f(\mathbb{B}^n) = \mathbb{B}^n\}$ then $\text{Möb}(\mathbb{B}^n)$ is generated by inversions in $(n - 1)$ -spheres orthogonal to $\mathbb{S}^{n-1} = \partial\mathbb{B}^n$ and is the isometry group of $\mathbb{B}^n$ with its hyperbolic metric. As before $\text{Möb}^+(\mathbb{B}^n) \subset \text{Möb}(\mathbb{B}^n)$ denotes those Möbius transformations which preserve orientation. The groups $\text{Möb}(\mathbb{B}^n)$ and $\text{Möb}^+(\mathbb{B}^n)$ preserve $\mathbb{S}^{n-1}$ and in the obvious way are isomorphic to $\text{Möb}(\mathbb{S}^{n-1})$ and $\text{Möb}^+(\mathbb{S}^{n-1})$ respectively.

Definition 2.1 *Let (X, d_X) and (Y, d_Y) be metric spaces and let $f : X \rightarrow Y$ be a map between them. Then if there is a constant $L > 0$ such that*

$$d_Y(f(x), f(y)) \leq L d_X(x, y)$$

for every $x, y \in X$, then f is L -Lipschitz. Furthermore, if we also have that

$$d_Y(f(x), f(y)) \geq d_X(x, y)/L$$

for every $x, y \in X$ then f is L -bilipschitz.

Definition 2.2 *If $f : (X, d_X) \rightarrow (Y, d_Y)$ is such that every $x \in X$ has a neighbourhood U such that the restriction of f to U is Lipschitz, then f is locally Lipschitz (similarly for the definition of locally bilipschitz)*

Remark 2.3 Metric spaces and locally Lipschitz maps form a category usually denoted by LIP. For this reason we will abbreviate locally Lipschitz to LIP. Moreover, if $f : (X, d_X) \rightarrow (Y, d_Y)$ is bijective and both f and f^{-1} are LIP maps, then we say f is a *lipeomorphism*.

We write $\text{LIP}(X)$ to denote the space of lipeomorphisms from X to itself. Since every LIP map with compact domain is Lipschitz, the space $\text{LIP}(\mathbb{S}^n)$ consists of all the bilipschitz homeomorphisms from $\mathbb{S}^n$ to itself (the point here is that we don't have to say *locally* bilipschitz). It is a standard result that every diffeomorphism from $\mathbb{S}^n$ to itself lies in $\text{LIP}(\mathbb{S}^n)$.

Remark 2.4 *We will only consider those homeomorphisms which preserve the orientation of $\mathbb{S}^n$.*

Let Ω, Ω' be subdomains of $\mathbb{R}^n$ and $f : \Omega \rightarrow \Omega'$ be a homeomorphism. Fix $x_0 \in \Omega$ and for small enough $\epsilon > 0$ consider the quantities

$$D_I(\epsilon) = \inf\{d_{\mathbb{R}^n}(f(x_0), y) : y \in f(\partial B_{\mathbb{R}^n}(x_0, \epsilon))\}$$

and

$$D_u(\epsilon) = \sup\{d_{\mathbb{R}^n}(f(x_0), y) : y \in f(\partial B_{\mathbb{R}^n}(x_0, \epsilon))\}$$

Let $K_{x_0}(\epsilon) = D_u(\epsilon)/D_l(\epsilon) \geq 1$. If there exists $K < \infty$ such that $K_{x_0} = \limsup_{\epsilon \rightarrow 0} K_{x_0}(\epsilon) < K$ for every $x_0 \in \Omega$ then we say f is K -quasiconformal. We say f is quasiconformal if f is K -quasiconformal for some K . If every point $x_0 \in \Omega$ has a neighbourhood U such that $f|_U$ is K -quasiconformal, then we say f is locally K -quasiconformal.

Locally quasiconformal homeomorphisms between open subsets of $\mathbb{R}^n$ form a pseudogroup on $\mathbb{R}^n$ (See [13] for the definition of a pseudogroup and more detail). In the usual way, this allows us to construct *quasiconformal manifolds*. These are manifolds whose transition maps are locally quasiconformal. Furthermore, if we have a homeomorphism $f : M \rightarrow N$ between two such manifolds, then we say it is locally quasiconformal if the following holds. For any pair of charts $\phi : M \rightarrow \mathbb{R}^n$ and $\psi : N \rightarrow \mathbb{R}^n$ the composition $\psi \circ f \circ \phi^{-1}$ is locally quasiconformal where defined. In particular, we let $\text{QC}(\mathbb{S}^n)$ denote the group of orientation preserving quasiconformal mappings from $\mathbb{S}^n$ to itself. Notice here that the standard smooth structure on $\mathbb{S}^n$ automatically induces a locally quasiconformal structure and as in the case of $\text{LIP}(\mathbb{S}^n)$, $\text{QC}(\mathbb{S}^n)$ contains all the orientation preserving diffeomorphisms of $\mathbb{S}^n$.

Remark 2.5 The space of lipeomorphisms between open subsets of $\mathbb{R}^n$ also forms a pseudogroup. So we could equally well have defined $\text{LIP}(\mathbb{S}^n)$ to be the group of orientation preserving bilipschitz homeomorphisms of $\mathbb{S}^n$ where its Lipschitz structure is induced from the standard smooth structure.

Throughout the rest of the paper we take $G(\mathbb{S}^n)$ to denote either $\text{LIP}(\mathbb{S}^n)$ or $\text{QC}(\mathbb{S}^n)$.

Take $\mathcal{B}$ to be the basis for the induced topology on $\mathbb{S}^n$ consisting of the open balls $\{B_{\mathbb{S}^n}(c, \epsilon) : c \in \mathbb{S}^n, \epsilon \in (0, \pi)\}$. This ensures that for all $B \in \mathcal{B}$, $\mathbb{S}^n \setminus B$ is homeomorphic to $\mathbb{B}^n$ by excluding the possibility that $B = \mathbb{S}^n$ or $\mathbb{S}^n$ minus a point. We do this to ensure that given two elements $B_1, B_2 \in \mathcal{B}$ we can find $g \in G(\mathbb{S}^n)$ such that $g(B_1) = B_2$.

If $g \in G(\mathbb{S}^n)$ is not the identity, then we say g is *supported* on $B \in \mathcal{B}$ if g is the identity outside B . Consequently, if $B_1 \subset B_2$ and $g \in G(\mathbb{S}^n)$ is supported on B_1 then g is also supported on B_2 . We set $G_0(\mathbb{S}^n)$ to be those elements of $G(\mathbb{S}^n)$ supported on elements of $\mathcal{B}$.

We have already mentioned that for all $B_1, B_2 \in \mathcal{B}$ there exists $g \in G(\mathbb{S}^n)$ with $g(B_1) = B_2$. In particular, if $B_1, B_2 \in \mathcal{B}$ and $g \in G_0(\mathbb{S}^n)$ is an element supported on B_1 , then g is conjugate to an element of $G_0(\mathbb{S}^n)$ supported on B_2 . Furthermore, we can choose g to lie in $\text{Möb}^+(\mathbb{S}^n) \subset G(\mathbb{S}^n)$ so that if $B \in \mathcal{B}$ then $g(B)$ will also lie in $\mathcal{B}$.

Definition 2.6 A pair $(\{B_i\}_{i \geq 0}, \rho)$ consisting of a sequence $\{B_i\}_{i \geq 0}$ of elements of $\mathcal{B}$, and $\rho \in G_0(\mathbb{S}^n)$ will be called a σ -*sequence* if,

1. $B_i \cap B_j = \emptyset$ if $i \neq j$
2. There exists $x_0 \in \mathbb{S}^n$ such that every neighbourhood of x_0 contains all but finitely many of the B_i
3. $\rho(B_i) = B_{i+1}$

This means that if ρ is supported on $B \in \mathcal{B}$ then $\bigcup_{i \geq 0} B_i \subset B$ (since condition 1 means that ρ cannot be the identity anywhere in $\bigcup_{i \geq 0} B_i$). If $(\{B_i\}_{i \geq 0}, \rho)$ is a sigma sequence, we will say it is supported on $B \in \mathcal{B}$ if ρ is.

Example 2.7 We now use the fact that $\text{Möb}^+(\mathbb{S}^n) \subset G(\mathbb{S}^n)$ to construct a σ -sequence explicitly. Consider $\mathbb{S}^n$ as $\mathbb{R}^n \cup \{\infty\}$ then the Möbius transformation,

$$f(\mathbf{x}) = \frac{\mathbf{x}}{r^2}, f(\infty) = \infty$$

with $r \in \mathbb{R} \setminus \{0\}$ is the time r^2 flow of the complete vector field,

$$\mathcal{X}(x_1, \dots, x_n) = -\sum_i x_i \frac{\partial}{\partial x_i}$$

Let $\varphi : \mathbb{S}^n \rightarrow \mathbb{R}$ be a smooth function which is identically one on $B_{\mathbb{S}^n}(0, 1)$ and zero outside $B_{\mathbb{S}^n}(0, 1 + \epsilon)$ for some $0 < \epsilon < 1$ and take ρ to be the time r^2 flow of $\varphi \cdot \mathcal{X}$. Then ρ will be a smooth map which is the same as f on $B_{\mathbb{S}^n}(0, 1)$ and the identity outside $B_{\mathbb{S}^n}(0, 1 + \epsilon)$ and so is in $G_0(\mathbb{S}^n)$. Now take $B \in \mathcal{B}$ with $B \subset B_{\mathbb{S}^n}(0, 1)$ and $\rho(B) \cap B = \emptyset$ and define $\{B_i\} = \rho^i(B)$ $i \geq 0$, these will all be in $\mathcal{B}$ since ρ is conformal on $B_{\mathbb{S}^n}(0, 1)$ and $(\{B_i\}, \rho)$ will be a σ -sequence supported on $B_{\mathbb{S}^n}(0, 1 + \epsilon)$ with $x_0 = 0$. Take any $B \in \mathcal{B}$, then since there exists $g \in G(\mathbb{S}^n)$ with $g(B_{\mathbb{S}^n}(0, 1)) = B$, we have that $(\{g(B_i)\}, g \circ \rho \circ g^{-1})$ is a σ -sequence supported on $g(B_{\mathbb{S}^n}(0, 1 + \epsilon))$.

Let $(\{B_i\}, \rho)$ be a σ -sequence supported on $B \in \mathcal{B}$, for each i take $g_i \in G_0(\mathbb{S}^n)$ which is supported on B_i . We would like to be able to say that $\prod_{i \geq 0} g_i = \dots \circ g_1 \circ g_0$ is also in $G_0(\mathbb{S}^n)$. If $g_i \in \text{QC}_0(\mathbb{S}^n)$ for each i and there exists $K \geq 1$ such that the g_i are all K -quasiconformal, then the composition $\prod_{i \geq 0} g_i$ is also K -quasiconformal and hence in $\text{QC}_0(\mathbb{S}^n)$. In particular, if $\rho|_{\cup B_i}$ is conformal (this was the case in the example we constructed above) and we take g_0 supported on B_0 which is K -quasiconformal. Then the conjugates $\rho^{-i} \circ g_0 \circ \rho^i$ will also be K -quasiconformal and supported on B_i . Hence, the composition $\prod_{i \geq 0} \rho^{-i} \circ g_0 \circ \rho^i$ will also be K -quasiconformal, and thus lie in $\text{QC}_0(\mathbb{S}^n)$.

Similarly, if for each i , $g_i \in \text{LIP}_0(\mathbb{S}^n)$ is supported on B_i and there exists $L \geq 1$ such that the Lipschitz constants of the g_i are bounded by L , then the composition $\prod_{i \geq 0} g_i$ is L -Lipschitz and hence in $\text{LIP}_0(\mathbb{S}^n)$. In particular, if we take g_0 supported on B_0 which is L -Lipschitz. Then again using ρ from the example above, the conjugates $\rho^{-i} \circ g_0 \circ \rho^i$ will also be L -Lipschitz and supported

on B_i . Hence, the composition $\prod_{i \geq 0} \rho^{-i} \circ g_0 \circ \rho^i$ will also be L -Lipschitz, and thus lie in $\text{LIP}_0(\mathbb{S}^n)$. We will use these observations to prove the following theorem.

Theorem 2.8 *Given a non-trivial $h \in G(\mathbb{S}^n)$, any element $g \in G_0(\mathbb{S}^n)$ is the product of four conjugates of h and h^{-1} .*

PROOF. As h is not the identity, we can find $B \in \mathcal{B}$ with $h^{-1}(B) \cap B = \emptyset$. Let $(\{B_i\}, \rho)$ denote a σ -sequence with $\cup_i B_i \subset B$ obtained as in Example 2.7, so that $\rho|_B$ is conformal (Note that ρ will be supported on some larger ball containing B). Since every element of $G_0(\mathbb{S}^n)$ can be conjugated in $G(\mathbb{S}^n)$ to one supported on B_0 it suffices to show that any g supported on B_0 is the product of four conjugates of h and h^{-1} .

So let $g \in G_0(\mathbb{S}^n)$ be supported on B_0 , then the map,

$$f = \prod_{n \geq 0} \rho^n \circ g \circ \rho^{-n}$$

is an element of $G_0(\mathbb{S}^n)$ supported on $\cup_{i \geq 0} B_i \subset B$. Since the conjugate $h^{-1} \circ f^{-1} \circ h$ is then supported on $h^{-1}(B)$ and $h^{-1}(B) \cap B = \emptyset$, we can think of the commutator,

$$f_1 := h^{-1} \circ f^{-1} \circ h \circ f = (h^{-1} \circ f^{-1} \circ h) \circ f$$

as two actions on disjoint sets.

The same can be said about,

$$f_2 := \rho \circ h^{-1} \circ f \circ h \circ \rho^{-1} = (\rho \circ h^{-1} \circ f \circ h \circ \rho^{-1}) \circ (\rho \circ f^{-1} \circ \rho^{-1})$$

here $\rho \circ f^{-1} \circ \rho^{-1}$ is supported on B whereas $\rho \circ h^{-1} \circ f \circ h \circ \rho^{-1}$ is supported on $h^{-1}(B)$.

So $f_2 \circ f_1$ can be examined in terms of its action on B and $h^{-1}(B)$ respectively. Doing this, one can easily see that $f_2 \circ f_1$ is the identity on $h^{-1}(B)$ and acts as g on B . Since g is supported on $B_0 \subset B$, $g = f_2 \circ f_1$, but we can write,

$$g = f_2 \circ f_1 = (\rho \circ h^{-1} \circ \rho^{-1}) \circ (\rho \circ f \circ h \circ f^{-1} \circ \rho^{-1}) \circ (h^{-1}) \circ (f^{-1} \circ h \circ f)$$

as a product of four conjugates of h and h^{-1} . □

Corollary 2.9 *The subgroup of $G(\mathbb{S}^n)$ generated by $G_0(\mathbb{S}^n)$ is simple.*

PROOF. Denote by $G(G_0(\mathbb{S}^n))$ the subgroup of $G(\mathbb{S}^n)$ generated by $G_0(\mathbb{S}^n)$ and let H be a non-trivial normal subgroup. We need to show that $H = G(G_0(\mathbb{S}^n))$. So let $g \in G(G_0(\mathbb{S}^n))$ and take $h \in H$ which is not the identity. We can write g as a finite composition $g_n \circ \dots \circ g_1$ where each $g_i \in G_0$ for $i = 1 \dots n$. Then using Theorem 2.8 we can write each g_i as a product of four conjugates of h and h^{-1} , but since H is normal this implies that each $g_i \in H$ and thus $g \in H$ which gives us the result. □

It now remains to show that $G(\mathbb{S}^n)$ is equal to $G(G_0(\mathbb{S}^n))$. We show this in the following section.

3 Sullivan groups

In this section we introduce the concept of a Sullivan group, this is a discrete group of hyperbolic isometries which gives rise to a hyperbolic n -manifold with the property that the complement of any point can be LIP-immersed into $\mathbb{R}^n$. We use this along with two versions of the famous Schoenflies Theorem in the LIP category to prove the main result, Lemma 3.6, which states that there is a neighbourhood of the identity in $G(\mathbb{S}^n)$ which is a subset of $G(G_0(\mathbb{S}^n))$. Using this, we prove that $G(G_0(\mathbb{S}^n))$ is equal to $G(\mathbb{S}^n)$ which then must be simple.

If (X, d_X) and (Y, d_Y) are metric spaces and $f : (X, d_X) \rightarrow (Y, d_Y)$ is a lipeomorphism onto its image (and hence injective) then we say f is a *LIP embedding* of X in Y . If every $x \in X$ has a neighbourhood U such that $f|_U : U \rightarrow Y$ is a LIP embedding then we say f is a *LIP immersion*.

Let $\Gamma \subset \text{Möb}^+(\mathbb{B}^n)$ be a discrete group of Möbius transformations which acts freely, then the quotient space $Q = \mathbb{B}^n/\Gamma$ is a smooth Hausdorff manifold and the natural map $\pi : \mathbb{B}^n \rightarrow Q$ is a covering map. Note that since Γ acts properly and $\mathbb{B}^n$ is locally compact, the property of Γ being discrete is equivalent to it acting properly discontinuously. Q inherits a hyperbolic metric from $\mathbb{B}^n$ defined in the usual way so that π becomes a local isometry and we write $B_Q(q, \epsilon)$ to denote the open ball of radius ϵ around $q \in Q$ in this metric.

Definition 3.1 *A group Γ of Möbius transformations acting on $\mathbb{B}^n$ is a **Sullivan Group** if it satisfies the following;*

1. Γ is discrete
2. Γ acts freely
3. $Q = \mathbb{B}^n/\Gamma$ is closed (compact and without boundary)
4. For every $q \in Q$ there exists a LIP immersion $\iota : Q \setminus q \rightarrow \mathbb{R}^n$

If Γ is a Sullivan group we call the quotient space $Q = \mathbb{B}^n/\Gamma$ a *Sullivan manifold*. We have the following theorem from [12].

Theorem 3.2 *There exists a Sullivan manifold of every dimension $n \geq 2$.*

Sullivan's proof of this theorem is deep and we take it on faith here. As an easy example though, in dimension 2 any closed surface of genus $g \geq 2$ can be constructed as a quotient of $\mathbb{B}^2$ by a group of Möbius transformations Γ which is discrete and acts freely. We claim that all of these groups are Sullivan groups.

To see this it remains to show condition 4, that for every $q \in Q$ there exists a LIP immersion $\iota : Q \setminus q \rightarrow \mathbb{R}^n$. So take $q \in Q$ and remove the ball $B_Q(q, \epsilon)$. The resulting surface is lipeomorphic (diffeomorphic in fact) to a disc with $2g$ strips attached (see Figure 1), which can then be LIP-immersed into




Figure 1: Immersion of genus 2 surface minus a disc into $\mathbb{R}^2$

$\mathbb{R}^n$. Letting ϵ tend to 0 then gives the required map (note however that the bilipschitz constants will become arbitrarily large near q).

We will also need the following two theorems which we state without proof.

Theorem 3.3 (Lipschitz Schoenflies Theorem [8]) *Suppose that $0 < a < 1$ and that f is a LIP embedding of the annulus $A = \bar{B}^n \setminus B_{\mathbb{R}^n}(0, a)$ into $\mathbb{S}^n$. Then $f|_{S^{n-1}}$ can be extended to a LIP embedding $f^* : \bar{B}^n \rightarrow \mathbb{S}^n$.*

Theorem 3.4 (Quantitative Schoenflies Theorem [15]) *Let $A = \bar{B}^n \setminus B_{\mathbb{R}^n}(0, 1/2)$, $\mathcal{N} = \{f : A \rightarrow \mathbb{R}^n : \sup\{d(x, f(x)) : x \in A\} < 1/30\}$ and $\mathcal{M} = \{f : \bar{B}^n \rightarrow \mathbb{R}^n\}$ then there exists a continuous map $\phi : \mathcal{N} \rightarrow \mathcal{M}$ a constant a_0 and constants b_n which depend only on n such that:*

1. $\phi(f)|_{S^{n-1}} = f|_{S^{n-1}}$ for all $f \in \mathcal{N}$
2. $\phi(id) = id$
3. If f is K -quasiconformal, then $\phi(f)$ is $a_n K^3$ -quasiconformal
4. If f is locally L -bilipschitz, $\phi(f)$ is locally $a_0 L^3$ -bilipschitz
5. If f is L -bilipschitz, $\phi(f)$ is $a_0 L^3$ -bilipschitz

Remark 3.5 *In the definition of $\mathcal{M}$ and $\mathcal{N}$ above we only consider $f : A \rightarrow \mathbb{R}^n$ which are embeddings.*

Lemma 3.6 *There exists $\epsilon > 0$ so that*

$$\{f \in G(\mathbb{S}^n) : \tilde{d}(id, f) < \epsilon\} \subset G(G_0(\mathbb{S}^n))$$

(Recall that $\tilde{d}(f, g) = \sup\{d_{\mathbb{S}^n}(f(x), g(x)) : x \in \mathbb{S}^n\}$).

PROOF. Take $f \in G(\mathbb{S}^n)$. Under the assumption that f is close to id we shall construct $g \in G_0(\mathbb{S}^n)$ which agrees with f on some open ball $B \in \mathcal{B}$. This will mean that $g^{-1} \circ f$ will be supported in the interior of B^c , and since $\text{int}(B^c) \in \mathcal{B}$, $g^{-1} \circ f$ will be in $G_0(\mathbb{S}^n)$. As a result $f = g \circ (g^{-1} \circ f)$ will be an element of $G(G_0(\mathbb{S}^n))$ as required and the proof will be complete.

Begin by using Theorem 3.2 to find a Sullivan group Γ , giving rise to a Sullivan manifold $Q = \mathbb{B}^n / \Gamma$ of dimension n . Let $\pi : \mathbb{B}^n \rightarrow Q$ denote the standard projection, then we have the following: For every $q \in Q$, there exists $r(q) > 0$ such that for any $p \in \pi^{-1}(q)$ there exists a neighborhood U of $\bar{B}_{\mathbb{B}^n}(r(q), p)$ such that the restriction $\pi|_U$ is an isometry. Since Q is compact we can fix $r > 0$ which works for every $q \in Q$ and define $D = B_{\mathbb{B}^n}(0, r) \subset \mathbb{B}^n$.

Consider the restriction of f to the closed unit ball $\bar{\mathbb{B}}^n \subset \mathbb{S}^n$, which we denote by f_0 . We shall construct $g : \mathbb{S}^n \rightarrow \mathbb{S}^n$ which is supported on $\mathbb{B}^n$ and agrees with f on D . It is thus necessary that we take f sufficiently close to the identity so as to ensure that $f_0(D) \subset \mathbb{B}^n$. Choose $q \in Q \setminus \pi(\bar{D})$ and use condition 4 of Definition 3.1 to find a LIP immersion $\iota : Q \setminus q \rightarrow B_{\mathbb{R}^n}(0, 0.9)$.

We can arrange that $\iota \circ \pi = id$ in a neighbourhood of $\bar{D}$. To see this, let U be a neighbourhood of $\bar{D}$ on which $\iota \circ \pi$ is a LIP-embedding. Let $B_1, B_2 \subset \mathbb{B}^n$ denote open balls centred at 0 which satisfy $\bar{D} \subset B_1 \subset \bar{B}_1 \subset B_2 \subset U$. Apply Theorem 3.3 to the map $(\iota \circ \pi)^{-1}$ on the annulus $B_2 \setminus B_1$ to find a LIP-embedding $h' : \mathbb{S}^n \setminus B_1 \rightarrow \mathbb{S}^n$ which agrees with $(\iota \circ \pi)^{-1}$ on ∂B_1 . Now define $h : \mathbb{S}^n \rightarrow \mathbb{S}^n$ by,

$$h(x) = \begin{cases} (\iota \circ \pi)^{-1}(x) & \text{if } x \in B_1 \\ h'(x) & \text{if } x \in \mathbb{S}^n \setminus B_1 \end{cases}$$

then h is a lipeomorphism which agrees with $(\iota \circ \pi)^{-1}$ in a neighbourhood of $\bar{D}$. By post-composing ι with h we will get a LIP immersion of Q into $\mathbb{S}^n$ whose image is contained in some element of $\mathcal{B}$ which contains $\bar{D}$. We may now use a diffeomorphism which is the identity on some neighbourhood of $\bar{D}$ to map $h(B_{\mathbb{R}^n}(0, 0.9))$ back inside $B_{\mathbb{R}^n}(0, 0.9)$. This will give us a new LIP immersion $\iota' : Q \setminus q \rightarrow B_{\mathbb{R}^n}(0, 0.9)$ with the additional property that $\iota' \circ \pi = id$ in a neighbourhood of $\bar{D}$. We now assume that $\iota \circ \pi = id$ in a neighbourhood of $\bar{D}$.

Choose $t > 0$ sufficiently small so that $3t < r$ and for $i = 1, 2, 3$ the balls $D_i = B_Q(q, it) \subset Q$, satisfy $D_i \cap \pi(\bar{D}) = \emptyset$.

We will construct g by lifting maps as shown.

$$\begin{array}{ccc}
g : & \mathbb{S}^n & \rightarrow \mathbb{S}^n \\
& \cup & \\
\tilde{f}_2 : & \mathbb{B}^n & \rightarrow \mathbb{B}^n \\
& \downarrow \pi & \downarrow \pi \\
f_2 : & Q & \rightarrow Q \\
& \cup & \\
f_1 : & Q \setminus D_2 & \rightarrow Q \setminus D_1 \\
& \downarrow \iota & \downarrow \iota \\
f_0 : & \mathbb{B}^n & \rightarrow \mathbb{S}^n
\end{array}$$

Firstly we construct $f_1 : Q \setminus D_2 \rightarrow Q \setminus D_1$, this is a natural lift of f_0 constructed using the LIP-immersion ι . In order to define f_1 we require that f is sufficiently close to id (i.e. that f_0 is close to the inclusion $\eta : \bar{B}^n \hookrightarrow \mathbb{R}^n \cup \{\infty\}$). This requirement is to ensure that $f_0 \circ \iota(Q \setminus D_2)$ is contained within the image of $\iota(Q \setminus D_1)$, a necessary condition for the map f_1 to exist.

To define f_1 , we first construct a finite open cover U_i for $i = 1, \dots, k$ of $Q \setminus D_2$, such that the restriction $\iota|_{U_i}$ is a LIP embedding for every i .

Now construct another finite cover W_i for $i = 1, \dots, k$, this time consisting of closed and hence compact subsets of $Q \setminus D_1$ such that $W_i \subset U_i$. By taking f closer to id if necessary we will have $f_0(\iota(W_i)) \subset \iota(U_i)$ so we can define f_1 on each W_i by $f_1|_{W_i} = (\iota|_{U_i})^{-1} f_0(\iota|_{W_i})$. This definition makes sense since ι is a LIP-embedding on each U_i and hence has a well defined inverse on its image.

Now by choosing f to lie in an even smaller neighbourhood of the identity if necessary, we can apply the quantitative Schoenflies theorem 3.4 to the restriction of f_1 to $\bar{D}_3 \setminus D_2$ to construct a map $f_2 : Q \rightarrow Q$ which agrees with f_1 on $Q \setminus D_3$. Now we can lift f_2 to $\tilde{f}_2 : \mathbb{B}^n \rightarrow \mathbb{B}^n$ in the usual way so that $\pi \circ \tilde{f} = f \circ \pi$. This lifting is unique if we specify $\tilde{f}_2(0) \in \pi^{-1} \circ f_2 \circ \pi(0)$, but we have only one choice since we want $\tilde{f}_2|_D = f_0|_D$, namely we set $\tilde{f}_2(0) = f_0(0)$.

Now, if we choose f to be still closer to the identity, then since $\iota(Q \setminus q)$ is compactly contained in $\mathbb{B}^n$, f_1 will move closer to the identity map of $Q \setminus D_2$ (Here we are using the hyperbolic metric on Q to determine this distance). Consequently the restriction of f_1 to ∂D_3 will also become closer to the identity (we used this to apply the quantitative Schoenflies theorem 3.4). Since the map ϕ in theorem 3.4 is continuous and it fixes id , $\phi(f_1|_{\bar{D}_3 \setminus D_2})$ will also move closer to the identity. As a result we are making f_2 close to the identity in the hyperbolic metric on Q .

If the dimension of Q is 3 or more then by taking f_2 close enough to the identity we can deduce that f_2 is homotopic to the identity by applying Mostow rigidity.

In dimension 2 however, we will show that f_2 will be homotopic to the

identity by showing that it acts as the identity on π_1 . To see this, take a set of generators $\tilde{\gamma}_1 \cdots \tilde{\gamma}_m$ for π_1 (this will be a finite set as Q is closed) and take a representative γ_i of each. Then for each γ_i there exists ϵ_i such that the ϵ_i -neighbourhood of γ_i is homotopy equivalent to γ_i . The same will then be true of $\epsilon' = \min\{\epsilon_i : i = 1, \dots, m\}$. One can now see that if f_2 is uniformly ϵ' close to the identity then $f_2(\gamma_i)$ will be homotopic to γ_i for each i . Consequently, it will induce the identity on π_1 .

The homotopy from f_2 to the identity then lifts to show that $\tilde{f}_2$ also homotopic to the identity. Now, since Q is compact there exists a compact region $K \subset \mathbb{B}^n$ such that $\pi(K) = Q$. Furthermore, the lifted homotopy commutes with the action of Γ and so is completely determined by its behaviour on K . Now, each point in K can only be moved a finite hyperbolic distance by the homotopy and hence the same applies to $\tilde{f}_2$. Since Γ acts by isometries this applies to every point in $\mathbb{B}^n$ allowing us to deduce that $\tilde{f}_2$ is a bounded hyperbolic distance from the identity.

We now use the fact that a lipeomorphism or K-quasiconformal homeomorphism of hyperbolic space which is a bounded hyperbolic distance from the identity will determine a map of the same class from $\bar{B}^n$ to itself which is the identity on $\partial\mathbb{B}^n$. This means that we can extend $\tilde{f}_2$ to a map $g : \mathbb{S}^n \rightarrow \mathbb{S}^n$ which agrees with $\tilde{f}_2$ on $\mathbb{B}^n$ and is the identity otherwise.

From the construction of g we have that $f_0\iota\pi = \iota\pi g$ where defined (in particular this applies in a neighbourhood of D) and $\iota\pi = id$ in a neighbourhood of D , consequently $f = g$ on D as we required.

It remains to check that g is in fact an element of $G_0(\mathbb{S}^n)$. Suppose that f was quasiconformal and let $E_1 = \pi^{-1}(\bar{D}_3)$ and $E_2 = \pi^{-1}(Q \setminus D_3)$. Then E_1 consists of countably many components E_1^i . Consider one such component E_1^1 , then the dilatations of $\tilde{f}_2$ on E_1^1 are determined by the behaviour of f_2 on $\bar{D}_3$. Since $f_2|_{\bar{D}_3}$ was defined by the quantitative Schoenflies theorem it is quasiconformal, and hence $\tilde{f}_2|_{E_1^1}$ will be too. Furthermore if $i \neq j$ there exists $\gamma \in \Gamma$ such that $\tilde{f}_2|_{E_1^i} = \gamma\tilde{f}_2\gamma^{-1}|_{E_1^j}$ since γ is conformal $g|_{E_1}$ is quasiconformal.

Now since Q is compact, there exists an open set $U \subset \mathbb{B}^n$ whose closure in $\mathbb{B}^n$ is compact and such that $\pi(U) = Q$, then $\iota\pi$ is locally bilipschitz on $U \setminus \pi^{-1}(\bar{D}_1)$. By construction, $\iota\pi\tilde{f}_2 = f_0\pi\iota$ in E_2 and hence $\tilde{f}_2|_{E_2 \cap U}$ is quasiconformal. Since $\gamma\tilde{f}_2 = \tilde{f}_2\gamma$ for all $\gamma \in \Gamma$, we get $\tilde{f}_2|_{E_2}$ is quasiconformal. It now follows (see [16] Theorem 35.1) that g is quasiconformal.

If f was a lipeomorphism of $\mathbb{S}^n$ then the same argument as above will allow us to deduce that $\tilde{f}_2$ is a lipeomorphism of $\mathbb{B}^n$ with its hyperbolic metric. This, along with the fact that $\tilde{f}_2$ is close to the identity, implies that $\tilde{f}_2$ is a lipeomorphism of $\mathbb{B}^n$ with the Euclidean metric and hence $g \in \text{LIP}(\mathbb{S}^n)$. $\square$

Theorem 3.7 $G(\mathbb{S}^n) = G(G_0(\mathbb{S}^n))$

PROOF. $G(G_0(\mathbb{S}^n))$ is a subgroup of the topological group $G(\mathbb{S}^n)$ which by Lemma 3.6 contains a neighbourhood of the identity and consequently we can translate this neighbourhood to show that $G(G_0(\mathbb{S}^n))$ is open in $G(\mathbb{S}^n)$. As $G(G_0(\mathbb{S}^n))$ is an open subgroup, it will be closed, as its cosets will also be open. Consequently, it suffices to show that $G(\mathbb{S}^n)$ is connected. So take $f \in G(\mathbb{S}^n)$, we can easily construct a path in $G(\mathbb{S}^n)$ which joins f to an element with a fixed point, so assume f has a fixed point $x \in \mathbb{S}^n$. Stereographic projection from x allows us to identify $\mathbb{S}^n \setminus \{x\}$ with $\mathbb{R}^n$. We can consequently think of f as a G -homeomorphism of $\mathbb{R}^n$, in particular there will exist $x_0 \in \mathbb{R}^n$ at which f is differentiable with positive Jacobian. Furthermore, we may assume that $x_0 = 0 = f(x_0)$. Set, $g_t(x) = f(tx)/t$ for $t > 0$ and $g_0 = f'(0)$, then the path g_t joins $f'(0)$ to f and since $\det(f'(0)) > 0$ we can join $f'(0)$ to the identity. We can extend all of these maps to $\mathbb{S}^n$ by leaving x invariant giving us a path in $G(\mathbb{S}^n)$ from f to the identity. This shows that $G(\mathbb{S}^n)$ is path connected and hence connected. □

References

- [1] R.D. Anderson, On Homeomorphisms as Products of Conjugates of a Given Homeomorphism and Its Inverse. *Topology of 3-Manifolds and Related Topics*, M. K. Fort, Ed., Prentice Hall, Englewood Cliffs, N.J., 1963, pp 231-234
- [2] A. Banyaga, Sur la structure du groupe des difféomorphismes qui préservent une forme symplectique. *Communicationes Mathematicae Helveticae*. 53, 1978, 174-227
- [3] M. Brown, H. Gluck, Stable Structures on Manifolds 1. Homeomorphisms of $\mathbb{S}^n$. *Ann. of Math.* 79, 1964, 1-17
- [4] M. Brown, H. Gluck, Stable Structures on Manifolds 2. Stable Manifolds. *Ann. of Math.* 79, 1964, 18-44
- [5] R.D. Edwards, R.C. Kirby, Deformations of Spaces of Imbeddings. *Ann. of Math.* 93, 1971, 63-88
- [6] D.B.A. Epstein, The Simplicity of Certain Groups of Homeomorphisms. *Composito Math.* 22, 1970, 165-173
- [7] R.C. Kirby, Stable homeomorphisms and the annulus conjecture, *Ann of Math.* 89, 1969, 575-582
- [8] J. Luukkainen, J. Väisälä, Elements of Lipschitz Topology. *Ann. Acad. Sci. Fenn. Ser. A I Math* 3, 1977, 85-122
- [9] J. Mather, Commutators of Diffeomorphisms. *Comment. Math. Helv.*, 49, 1974, 512-528
- [10] J. Mather, Commutators of Diffeomorphisms II. *Comment. Math. Helv.*, 50, 1975, 33-40
- [11] D. McDuff, On the group of volume preserving diffeomorphisms of $\mathbb{R}^n$. *Trans. Amer. Math. Soc.* 261, 1980, 103-113
- [12] D. Sullivan, Hyperbolic geometry and homeomorphisms. *Geometric Topology* (Proceedings Georgia Topology Conf., Athens, Ga., 1977) edited by J.C. Cantrell, Academic Press, New York, N.Y. London, 1979, 543-555
- [13] W.P. Thurston, *Three-Dimensional Geometry and Topology*. Volume 1. Princeton University Press, Princeton, New Jersey. 1997.
- [14] W.P. Thurston, Foliations and Groups of Diffeomorphisms. *Bull. Amer. Math. Soc.*, 80, 1974, 304-307
- [15] P. Tukia, J. Väisälä, Lipschitz and Quasiconformal approximation and extension. *Ann. Acad. Sci. Fenn. Ser. A I Math* 6, 1981, 303-342

- [16] J. Väisälä, Lectures on n -dimensional quasiconformal mappings. Lecture Notes in Mathematics 229, Springer-Verlag, Berlin-Heidelberg-New York, N. Y., 1971.